



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA®
FOR CONTROL OF
BLUE MOLD
ON TOBACCO

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®
EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Tobacco	Blue Mold (<i>Peronospora tabacina</i>)	Apply Regalia® preventatively at a rate of 1% v/v in 50-100 gallons of water per acre.	Apply Regalia® preventatively at a rate of 1% v/v in 50-100 gallons of water per acre. For improved performance, use Regalia in a tank mix or rotational program with other registered fungicides.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2010

Marrone Bio Innovations, Inc.
2121 Second Street, Suite B-107
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com