



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA®
TO CONTROL PYTHIUM (AERIAL BLIGHT PHASE)
ON LEGUMES

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®
EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Legumes/Vegetables such as Beans, Green Beans, Snap Beans, Shell Beans, Dry Beans, Garbanzo Beans, Lima Beans, Peas, Chick Peas, Split Peas, and Lentils.	Pythium (Aerial Blight Phase) (<i>Pythium spp.</i>)	Apply 2-4 quarts Regalia per acre when used alone. For improved performance, use Regalia at 1-4 quarts per acre in a tank mix or rotational program with another registered fungicide. Apply Regalia preventatively in 20-100 gallons of water per acre.	Begin applications preventatively (before disease symptoms become visible). Repeat applications at 7 to 14 day intervals as needed.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
2121 Second Street, Suite B-107
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com