



TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF REGALIA®  
TO CONTROL LATE BLIGHT  
ON CELERY

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**REGALIA®**  
**EPA Reg. No. 84059-3**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Celery	Late Blight ( <i>Septoria apiicola</i> )	Regalia is recommended to be used at 1-4 quarts per acre in a tank mix or rotational program with another registered fungicide.  Regalia may be used alone at 2-4 quarts per acre.  Apply Regalia preventatively in 50-200 gallons of water per acre.	Begin applications preventatively (before disease symptoms become visible).  Repeat applications at 7 to 14 day intervals as needed.

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.  
© Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.  
2121 Second Street, Suite B-107  
Davis, California 95618  
Telephone (530) 750-2800  
www.marronebioinnovations.com

**2121 Second Street, Suite B-107 • Davis, CA 95618 • Phone: 530-750-2800**