



TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF REGALIA®  
AS A PRE-PLANT DIP  
ON STRAWBERRY

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**REGALIA®**  
**EPA Reg. No. 84059-3**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

CROP	DISEASES	RATE	USE RECOMMENDATIONS
Strawberry	Black root rot ( <i>Rhizoctonia</i> spp., <i>Pythium</i> spp., <i>Fusarium</i> spp., <i>Cylindrocarpon</i> spp.)  Colletotrichum crown rot ( <i>Colletotrichum</i> spp.)  Phytophthora root rot and crown rot ( <i>Phytophthora</i> spp.)  Verticillium wilt ( <i>Verticillium</i> spp.)	Recommended use rate is a 0.5% to 1.0% suspension of Regalia (1 gallon Regalia in 100 to 200 gallons of water).	Apply as a pre-plant dip to strawberry plants, roots and crowns for 5 minutes. Treat transplants immediately prior to planting.  Regalia is compatible with other root dip treatments.

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.  
© Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.  
2121 Second Street, Suite B-107  
Davis, California 95618  
Telephone (530) 750-2800  
www.marronebioinnovations.com

**2121 Second Street, Suite B-107 • Davis, CA 95618 • Phone: 530-750-2800**