



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® TO CONTROL
ANTHRACNOSE FRUIT ROT
AND ALTERNARIA FRUIT ROT
ON BERRIES

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®
EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Berries such as Blueberry, Blackberry, Cranberry, Elderberry, Gooseberry, Loganberry, Raspberry, and Other Berry Crops	Anthracnose Fruit Rot (<i>Colletotrichum acutatum</i>) Alternaria Fruit Rot (<i>Alternaria spp.</i>)	Apply 1-2 quarts of Regalia® in 50-100 gallons of water per acre.	For control of Anthracnose Fruit Rot and Alternaria Fruit Rot, apply 1-2 quarts of Regalia® in 50-100 gallons of water per acre. Initiate application at green tip and continue applications on 7- to 10-day intervals until 10% blue fruit (blueberry) or 10% ripe fruit for other berry crops. Regalia® may be applied up to and including the day of harvest (0-day PHI).

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
2121 Second Street, Suite B-107
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com